

# **Analysis Report** Honey-Profiling TM

Sample ID: PI2407150255

# Information/Declaration provided by customer:

Type of Sample: Honey
Type of Honey: Unknown
Botanical Variety: undefined
Geographical Origin: India

Disclaimer: this information will affect the applicability and validity of analyses and results.

Note: it is important to fill in these information in a correct and precise manner (e.g. variety in case of monofloral honey, and country of origin). The tests applied (and therefore the results received) are different from one type of honey to the other. Bruker does not take responsibility for wrong or incomplete information given by the customer.

Measuring Date: 17-Jul-2024 14:53:33

Reporting Date: 17-Jul-2024 15:02:15, 13 pages, Version 3.1.3

#### **Results Summary**

Type of Analysis	Result	Status
Origin, Type and Variety		
Type of Honey	Blossom	
Botanical Variety	Not Applied	
Country of Origin: India	Not Consistent	
Test of Markers of Foreign Sugars	Pass	
Composition and Comparison to Reference Group	Requires Interpretation	
Compliance for EU Market for		
HMF	Compliant	
Sucrose	Compliant	
Glucose + Fructose	Compliant	

The data analysis is performed at Bruker BioSpin GmbH (Ettlingen, Germany) according to testing method AA-72-03-18 (Honey-Profiling 3.1.3), released on 09-Nov-2023 (DIN EN ISO/IEC 17025:2018 Accreditation Certificate D-PL-19229-01-00). All results solely refer to the tested sample as provided by the customer.




# **Origin, Type and Variety**

# **Type of Honey**

(Analysis-ID: HS3-CC-1000-18151)

This model is based on 14146 samples, thereof 1993 samples of reference group *Honeydew*.

**Result:** Consistent with *Blossom*. The probability of consistency is 100.0%.



## **Botanical Variety**


Verification of consistency is not possible as the declared botanical variety of the sample is unknown or Polyfloral.

# Country of Origin: India

(Analysis-ID: HS3-CC-116-74243)

This model is based on 25067 samples, thereof 744 samples of reference group *India*.

**Result:** Not consistent with declared country *India*. The probability of consistency is 0.0%.





# Test of Markers of Foreign Sugars

(Analysis-ID: HO-TAD-01/0833)

Following tests have been applied in order to detect foreign sugars:

ID	Description	Value	Graphics	Result
2	3.8113 / (1.95-2.02)	12.38	· // //	pass
3	3.9238 / (3.16-4.14)	0.0004819	Maria and a	pass
4	3.2890 / (1.95-2.02)	1.431	•	pass
5	4.0363 / (3.75-3.80)	0.00643		pass
6	5.3324 / (5.27-5.30)	0.0660		pass
7	3.0123 / (1.95-2.02)	0.01378	•	pass
8	3.5929	1694	•	pass
9	5.0745 / (4.07-4.12)	0.003556	Maria de la companione	pass
10	3.2753	231.2	•	pass
11	3.7490 / (1.95-2.02)	0.14	•	pass
13	4.1961 / (3.16-4.14)	0.0001079	•	pass
14	3.3135 / (5.27-5.30)	0.0197	•	pass
15	4.6809 / (3.65-3.73)	0.00013698		pass
16	3.7715 / (3.65-3.73)	0.04890	•	pass
17	5.4913 / (4.60-4.67)	0.001919	•	pass
18	4.6327 / (3.16-4.14)	0.0004809	Maria de la companione	pass
19	4.2569 / (1.95-2.02)	0.0292	•	pass
20	3.2797	165.0	•	pass
127	4.2487	133.6	•	pass
128	5.3045 / (1.95-2.02)	0.1024	•	pass
129	3.5884 / (3.75-3.80)	0.004975	•	pass
130	4.9421 / (4.93-4.97)	0.1522	· ·	pass
131	3.5929 / (4.07-4.12)	0.01225	•	pass
133	4.9433 / (4.93-4.97)	0.07545	•	pass
134	4.2501	55.4	•	pass
135	3.9156 / (4.07-4.12)	0.02863	<b>/</b>	pass
136	5.2876 / (4.07-4.12)	0.002688	•	pass
140	3.2852 / (4.07-4.12)	0.006249	· Mariana de la compansión de la compans	pass
142	4.3176 / (5.27-5.30)	0.02001		pass
143	5.3371 / (5.27-5.30)	0.01713	• <mark>//</mark>	pass
144	4.4347 / (4.07-4.12)	0.0002241	// <sub></sub>	pass
155	4.1511	447.1		pass
156	3.3701 / (3.20-3.22)	0.08911	/ <mark>/</mark>	pass
157	4.1781 / (4.07-4.12)	0.0008987	// <sub>_</sub>	pass
158	3.9283 / (5.27-5.30)	0.0750	·	pass



ID	Description	Value	Graphics	Result
159	3.9246 / (5.20-5.25)	0.02227	Marian Parante	pass
162	3.1784 / (5.20-5.25)	0.004483	A Committee of the Comm	pass
163	3.5929 / (3.16-4.14)	0.0006181	• · · · · · · · · · · · · · · · · · · ·	pass
166	3.6627 / (4.07-4.12)	0.03577	Marian and a	pass
167	3.2977 / (4.07-4.12)	0.003099	// <sub></sub>	pass
168	3.3753 / (4.60-4.67)	0.1687	•	pass
169	3.7579 / (3.65-3.73)	0.12001	•	pass
170	3.8885 / (3.75-3.80)	0.04902	Maria de la companya della companya della companya della companya de la companya della companya	pass
171	4.1651 / (1.95-2.02)	1.502	• • • • • • • • • • • • • • • • • • •	pass
172	3.1637	19.307	//	pass
191	3.7932 / (4.07-4.12)	0.2373	· · · · · · · · · · · · · · · · · · ·	pass
1000	min. fructose/glucose	1.47	Maria and an and	pass
1001	max. fructose/glucose	1.47		pass
1002	turanose [g/100g]	2.03	M	pass
1003	sucrose [g/100g]	1.4	• · · · · · · · · · · · · · · · · · · ·	pass
1004	total sugar [g/100g]	72.6	Maria and a second	pass
1005	proline [mg/kg]	636	Maria and the	pass
1006	DHA(D), mannose(M) [mg/kg]	D=0.0, M=39		pass

Result: There are no indications for the presence of foreign sugars.

#### Notes:

- The column *ID* is the marker's identification number.
- The column *Description* indicates either the NMR spectral region(s) (in ppm) concerned by the marker, or the molecule when it has been identified.
- The column Value is the result obtained for the marker.
- For columns *Graphics* and *Result* please refer to detailed description in section "General Remarks" at the end of this report.



#### **DHA** and Mannose

Compound	Value	Unit	LOQ	Reference Range	Flag
mannose	<loq< td=""><td>g/100g</td><td>0.02</td><td>&lt;0.02 g/100g in reference dataset</td><td></td></loq<>	g/100g	0.02	<0.02 g/100g in reference dataset	
dihydroxyacetone (DHA)	<loq< td=""><td>mg/kg</td><td>5</td><td>&lt; 5</td><td></td></loq<>	mg/kg	5	< 5	

Note: the reference range is based on 327 India samples in the Honey-Profiling Database.

#### **Guideline:**

- Mannose is a mono saccharide not found in honeys with a pH value lower than 5, but that is regularly found in industrial sugars. In rare cases, however, the presence of mannose cannot be excluded for certain geographic origins and/or botanical varieties. A concentration of mannose exceeding 0.02 g/100g in honey with a pH < 5 could indicate the presence of foreign sugars or industrial processing practices which are not suitable for honey. An expert interpretation is suggested when mannose is present in the honey.</p>
- Dihydroxyacetone and/or methylglyoxal are only known to be naturally present in Leptospermum genus honeys from Australia and New Zealand. A concentration exceeding 5 mg/kg in other types of honey is not typical and could indicate the presence of foreign sugars or industrial processing practices which are not suitable for honey. An expert interpretation is suggested in such cases.
- DHA and mannose are often observed simultaneously in various types of sugar syrups.

Deviations in the sugar profile, fermentation parameters and comparison to reference group could also indicate adulterations. Please check the section "Composition and Comparison to reference group" in addition.



# **Composition and Comparison to Reference Group**

# **Quantitative Analysis of Compounds**

(Analysis-ID: HO-Q/3.1.0)

Parameters labelled with  $^*$  are calculated parameters. The reference range is based on 327 India samples in the Honey-Profiling Database.

### Sugars:

Compound	Value	Unit	LOQ	Reference Range	Flag
glucose + fructose *	72.6	g/100g	20.0	60.0 77.2	
fructose / glucose *	1.47	_	_	0.96	
fructose	43.2	g/100g	10.0	31.4 42.2	
glucose	29.4	g/100g	10.0	26.9 38.6	
sucrose	1.4	g/100g	0.5	< 0.5 g/100g in reference dataset	
turanose	2.03	g/100g	0.2	0.5	
maltose	2.9	g/100g	0.5	<0.5	
melezitose	<loq< td=""><td>g/100g</td><td>1.0</td><td>&lt;1.0 g/100g in reference dataset</td><td></td></loq<>	g/100g	1.0	<1.0 g/100g in reference dataset	
maltotriose	<loq< td=""><td>g/100g</td><td>1.0</td><td>&lt;1.0 g<math>/</math>100g in reference dataset</td><td></td></loq<>	g/100g	1.0	<1.0 g $/$ 100g in reference dataset	
gentiobiose	<loq< td=""><td>g/100g</td><td>0.3</td><td>&lt;0.3 g/100g in reference dataset</td><td></td></loq<>	g/100g	0.3	<0.3 g/100g in reference dataset	
raffinose	0.1	g/100g	0.1	0.1 0.3	

#### Acids:

Compound	Value	Unit	LOQ	Reference Range	Flag
citric acid	86	mg/kg	50	<50 162	
malic acid	<loq< td=""><td>mg/kg</td><td>100</td><td>&lt;100 115</td><td></td></loq<>	mg/kg	100	<100 115	
quinic acid	<loq< td=""><td>mg/kg</td><td>300</td><td>&lt;300 mg/kg in reference dataset</td><td></td></loq<>	mg/kg	300	<300 mg/kg in reference dataset	



# Amino Acids:

Compound	Value	Unit	LOQ	Reference Range	Flag
alanine	13	mg/kg	5	<5 24	
aspartic acid	<loq< td=""><td>mg/kg</td><td>150</td><td>&lt;150 mg/kg in reference dataset</td><td></td></loq<>	mg/kg	150	<150 mg/kg in reference dataset	
glutamine	<loq< td=""><td>mg/kg</td><td>200</td><td>&lt;200 mg/kg in reference dataset</td><td></td></loq<>	mg/kg	200	<200 mg/kg in reference dataset	
leucine	<loq< td=""><td>mg/kg</td><td>40</td><td>&lt;40 110</td><td></td></loq<>	mg/kg	40	<40 110	
proline	636	mg/kg	200	225 863	
valine	<loq< td=""><td>mg/kg</td><td>10</td><td>&lt;10 14</td><td></td></loq<>	mg/kg	10	<10 14	
tyrosine	<loq< td=""><td>mg/kg</td><td>50</td><td>&lt;50 150</td><td></td></loq<>	mg/kg	50	<50 150	
phenylalanine	<loq< td=""><td>mg/kg</td><td>100</td><td>&lt;100 568</td><td></td></loq<>	mg/kg	100	<100 568	

# Indicators for Fermentation and Processing:

Compound	Value	Unit	LOQ	Reference Range	Flag
2,3-butanediol	<loq< td=""><td>mg/kg</td><td>20</td><td>&lt;20 327</td><td></td></loq<>	mg/kg	20	<20 327	
5-hydroxymethylfurfural (HMF)	<loq< td=""><td>mg/kg</td><td>5</td><td>&lt;5 122</td><td></td></loq<>	mg/kg	5	<5 122	
acetic acid	17	mg/kg	10	<10 113	
acetoin	<loq< td=""><td>mg/kg</td><td>20</td><td>&lt;20 187</td><td></td></loq<>	mg/kg	20	<20 187	
ethanol	<loq< td=""><td>mg/kg</td><td>5</td><td>&lt;5 3283</td><td></td></loq<>	mg/kg	5	<5 3283	
lactic acid	21	mg/kg	10	<10 234	
formic acid	65	mg/kg	5	<5 113	
fumaric acid	<loq< td=""><td>mg/kg</td><td>5</td><td>&lt;5 7</td><td></td></loq<>	mg/kg	5	<5 7	
pyruvic acid	17	mg/kg	10	<10 32	
succinic acid	11	mg/kg	5	8 94	



#### Markers:

Compound	Value	Unit	LOQ	Reference Range	Flag
3-phenyllactic acid	<loq< td=""><td>mg/kg</td><td>300</td><td>&lt;300 mg/kg in reference dataset</td><td></td></loq<>	mg/kg	300	<300 mg/kg in reference dataset	
methylglyoxal (MGO)	<loq< td=""><td>mg/kg</td><td>30</td><td>&lt;30 33</td><td></td></loq<>	mg/kg	30	<30 33	
kynurenic acid	<loq< td=""><td>mg/kg</td><td>60</td><td>&lt;60 mg/kg in reference dataset</td><td></td></loq<>	mg/kg	60	<60 mg/kg in reference dataset	
shikimic acid	<loq< td=""><td>mg/kg</td><td>80</td><td>&lt;80 mg/kg in reference dataset</td><td></td></loq<>	mg/kg	80	<80 mg/kg in reference dataset	

#### **Guideline:**

- Values of fructose/glucose ratio exceeding 1.95 or below 0.85 are not typical for honey and could indicate the presence of sugar syrups.
- Low concentrations of turanose (less than 0.35 g/100g) is a marker for the presence of foreign sugars (see section "Markers of Foreign Sugars").
- Atypical concentrations of sugars in comparison to reference range can be related to foreign sugars/adulteration or to the botanical varieties present in the honey. Expert interpretation is suggested.
- The presence of gentiobiose is common for Linden Tree honey.
- The presence of quinic acid is common for honeydew.
- HMF is a sugar degradation product formed under influence of heat. According to EU-Directive (2001/110/EC), HMF can be found naturally in honey in concentrations up to 80 mg/kg in regions with tropical climate. A higher concentration of HMF is due to processing of honey which can sometime correlate with adulterations. A concentration of HMF exceeding 200 mg/kg should be regarded as suspicious.
- Concentration of ethanol exceeding 400 mg/kg indicates fermentation of the honey, which could be related to unripe honey.
- Acetoin can be elevated in Eucalyptus honey.
- The presence of kynurenic acid is common for Chestnut honey.
- The presence of shikimic acid is common for honeydew.



#### Statistical Comparison with the Reference Group

The models are based on 327 samples from *India* in the Honey-Profiling Database.

#### **Univariate Verification**

(Analysis-ID: HS3-NTV-116-74261)

Applied Model: India

Result: Not typical for India.

 $3.591^{high}\ 4.186^{high}\ 4.208^{high}\ 5.379^{high}\ 5.383^{high}\ 5.388^{high}\ 5.392^{high}\ 5.397^{high}\ 5.406^{high}\ 5.599^{high}\ 5.604^{high}$ 

The suffix *high* indicates that the intensity is higher than in the reference group, the suffix *low* indicates that the intensity is lower than in the reference group.

**Possible molecules:** 2,3-Butanediol, Fructose, Gentiobiose, Maltotriose, Mannose, Melezitose, Proline, Quinic acid, Raffinose, Sucrose, Turanose

This list gives indication about possible molecules responsible for deviations.

#### **Multivariate Verification**

(Analysis-ID: HS3-NTV-116-74261)

**Applied Model:** India **Result:** Typical for India.

# Off-Model In-Model Amount of the control of the co

log-Mahalanobis Parameter



# Codex Alimentarius and EU-Directive 2001/110/EC:

Following parameters are required according to Codex Alimentarius and EU-Directive 2001/110/EC. The concentrations are obtained by direct quantification. Parameters labelled with  $^{\ast}$  are calculated parameters.

				Official Reference		
Compound	Value	Unit	LOQ	min	max	Flag
glucose + fructose *	72.6	g/100g	20.0	45	-	
sucrose	1.4	g/100g	0.5	_	15	
5-hydroxymethylfurfural (HMF)	<loq< td=""><td>mg/kg</td><td>5</td><td>_</td><td>80</td><td></td></loq<>	mg/kg	5	_	80	

Following flags are used according to Codex Alimentarius and EU-Directive 2001/110/EC:

Compound	Flag	Concentration	Declaration	Interpretation
glucose +		< 45 g/100g	All	Not compliant
fructose		< 60 g/100g	Blossom	Not compliant for blossom honey
		$\geq$ 60 g/100g	All	Compliant
		$\geq$ 45 g/100g	Honeydew	Compliant for honeydew honey
		$\geq$ 45 g/100g, $<$ 60 g/100g	Unknown	Compliant for honeydew honey and blends of honeydew honey with blossom honey. Not compliant for blossom honey
sucrose		> 15 g/100g	All	Not compliant
		10-15 g/100g	Acacia, Eucalyptus	Not compliant for false acacia (Robinia pseudoacacia), and red gum (Eucalyptus camadulensis)
		$\leq$ 5 g/100g	All	Compliant
		$\leq$ 10 g/100g	Acacia, Eucalyptus	Compliant for false acacia ( <i>Robinia pseu doacacia</i> ), and red gum ( <i>Eucalyptus ca madulensis</i> )
		$\leq$ 15 g/100g	Lavender	Compliant for Lavandula spp.
		5-10 g/100g	All, except Acacia, Eucalyptus, Lavender	If $\leq 15 \mathrm{g}/100 \mathrm{g}$ : compliant for lavende (Lavandula spp.) and borage (Borago of ficinalis). If $\leq 10 \mathrm{g}/100 \mathrm{g}$ : compliant fo false acacia (Robinia pseudoacacia), alfalfa (Medicago sativa), Menzies Banksia (Banksia menziesii), French honeysuckle (Hedysarum), red gum (Eucalyptus camadulensis), leatherwood (Eucryphia lucida, Eucryphia milliganii) and Citrus spp
HMF		$> 80~{ m mg/kg}$	All, except Indus- trial honey	Not compliant, except for baker's honey
		$\leq$ 40 mg/kg	All	Compliant
		> 80 mg/kg	Industrial honey	Compliant for baker's honey
		40-80 mg/kg	All	Not compliant, except for baker's honey and honeys of declared origin from regions with tropical climate and blends of these honeys



#### **General Remarks**

#### **Targeted Markers for Foreign Sugars**

This test aims at detecting foreign sugars. It does rely on the analysis of the concentrations of certain known molecules and on absolute intensities or intensities ratios of marker peaks. The Honey-Profiling report contains several tens of markers. Some of these markers are independent on the variety and the origin, in order to be able to analyze blends, while there are also many markers specific to countries or varieties.

The values obtained for each marker are compared to the thresholds for purity.

Reported and visualized results:

#### • Result Pass (green):

Markers with an upper limit: The value of the marker is below the acceptance limit Markers with a lower limit: The value of the marker is above the acceptance limit Conclusion: There are no indications for the presence of foreign sugars.

#### • Result Conditional Pass (yellow):

The value of the marker is between the acceptance limit and the threshold Conclusion: There is no indication for the presence of foreign sugars. However, for one or more markers the determined values are close to the threshold.


#### • Result Conditional Fail: (orange)

The value of the marker is between the threshold and the tolerance limit Conclusion: There is a strong indication for the presence of foreign sugars. However, the determined values are close to the threshold. It is recommended to perform additional tests in order to better evaluate the risk.

#### • Result Fail: (red)

Markers with an upper limit: The value of the marker is above the tolerance limit Markers with a lower limit: The value of the marker is below the tolerance limit Conclusion: There is a strong indication for the presence of foreign sugars.

The black dot represents the current sample.



Presence of foreign sugars is due to deliberate dilution of honey with sugar syrups or wrong beekeeping practices in terms of artificial feeding of bees.

All conformity statements in this section are based on the "Non Binary Statement with Guard Band" according to ILAC G8:09/2019. Please find in the following table the values for acceptance limit (conditional pass), threshold (conditional fail) and tolerance limit (fail).

ID	Description	Conditional Pass	Conditional Fail	Fail
2	3.8113 / (1.95-2.02)	>68.95	>69.44	>69.92
3	3.9238 / (3.16-4.14)	< 0.0003172	< 0.0003039	< 0.0002905
4	3.2890 / (1.95-2.02)	>5.792	>5.838	>5.884
5	4.0363 / (3.75-3.80)	>0.01113	>0.01131	>0.01148
6	5.3324 / (5.27-5.30)	>0.1683	>0.1703	>0.1723
7	3.0123 / (1.95-2.02)	>0.06929	>0.07134	>0.07340



ID	Description	Conditional Pass	Conditional Fail	Fail
8	3.5929	>2186	>2227	>2269
9	5.0745 / (4.07-4.12)	< 0.001662	< 0.001495	< 0.001329
10	3.2753	>407.8	>414.9	>422.0
11	3.7490 / (1.95-2.02)	>11.53	>11.56	>11.59
13	4.1961 / (3.16-4.14)	>0.0002284	>0.0002327	>0.0002369
14	3.3135 / (5.27-5.30)	>0.1677	>0.1717	>0.1757
15	4.6809 / (3.65-3.73)	< 0.00009264	< 0.00008224	< 0.00007184
16	3.7715 / (3.65-3.73)	>0.06283	>0.06399	>0.06515
17	5.4913 / (4.60-4.67)	>0.002301	>0.002341	>0.002380
18	4.6327 / (3.16-4.14)	< 0.0003476	< 0.0003379	< 0.0003282
19	4.2569 / (1.95-2.02)	>0.7715	>0.7759	>0.7803
20	3.2797	>262.1	>268.0	>273.9
127	4.2487	>366.4	>382.3	>398.2
128	5.3045 / (1.95-2.02)	>0.6017	>0.6092	>0.6168
129	3.5884 / (3.75-3.80)	>0.006261	>0.006470	>0.006679
130	4.9421 / (4.93-4.97)	>0.1832	>0.1862	>0.1892
131	3.5929 / (4.07-4.12)	>0.01423	>0.01446	>0.01470
133	4.9433 / (4.93-4.97)	>0.08613	>0.08758	>0.08904
134	4.2501	>237.5	>244.2	>251.0
135	3.9156 / (4.07-4.12)	< 0.02216	< 0.02141	< 0.02065
136	5.2876 / (4.07-4.12)	>0.004380	>0.004442	>0.004505
140	3.2852 / (4.07-4.12)	>0.008687	>0.008959	>0.009231
142	4.3176 / (5.27-5.30)	>0.07307	>0.07400	>0.07494
143	5.3371 / (5.27-5.30)	>0.04644	>0.04885	>0.05127
144	4.4347 / (4.07-4.12)	< 0.0001627	< 0.0001543	< 0.0001460
155	4.1511	<307.6	<268.2	<228.8
156	3.3701 / (3.20-3.22)	< 0.07742	< 0.07561	< 0.07380
157	4.1781 / (4.07-4.12)	< 0.0004571	< 0.0003843	< 0.0003115
158	3.9283 / (5.27-5.30)	>0.4156	>0.4226	>0.4295
159	3.9246 / (5.20-5.25)	< 0.01303	< 0.01167	< 0.01031
162	3.1784 / (5.20-5.25)	< 0.002753	< 0.002590	< 0.002427
163	3.5929 / (3.16-4.14)	>0.0008186	>0.0008317	>0.0008449
166	3.6627 / (4.07-4.12)	< 0.02052	< 0.02000	< 0.01949
167	3.2977 / (4.07-4.12)	< 0.002394	< 0.002260	< 0.002126
168	3.3753 / (4.60-4.67)	>0.1892	>0.1935	>0.1977
169	3.7579 / (3.65-3.73)	< 0.08739	< 0.08315	< 0.07891
170	3.8885 / (3.75-3.80)	< 0.03880	< 0.03725	< 0.03570
171	4.1651 / (1.95-2.02)	>6.362	>6.421	>6.480
172	3.1637	<10.563	< 8.554	< 6.544
191	3.7932 / (4.07-4.12)	>0.3806	>0.3999	>0.4192
1000	min. fructose/glucose	< 0.90	< 0.85	< 0.80
1001	max. fructose/glucose	>1.90	>1.95	>2.00
1002	turanose [g/100g]	< 0.38	< 0.35	< 0.32
1003	sucrose [g/100g]	>14	>15	>16
1004	total sugar [g/100g]	<46	<45	<44
1005	proline [mg/kg]	<171	<160	<149
1006	DHA(D), mannose(M) [mg/kg]	D>3, M>150	D>5, M>200	D>7, M>250

#### **Classification Models**

Analysis of origin, type and variety relies on a statistical classification analysis. The test applied is a classification analysis with the aim to check the consistency of the declared meta-information of the sample (geographical origin or botanical variety). The consistency with a group is expressed as posterior probability in the range from 0% to 100%. A posterior probability exceeding 50% is being regarded as consistent with the respective group. The underlying statistical models are based on a dimension reduction (Principal Component Analysis and/or Linear Discriminant Analysis used) followed by a Linear (or Quadratic) Discriminant Analysis for final classification.

Within the discrimination space figure, the ellipsoids are representing the modeling samples and the star represents the actual sample under investigation.



Verification of origin is not possible on blends from different countries.

Expert interpretation is necessary before deducing any conclusions.

All conformity statements in this section are based on the "Binary Statement for Simple Acceptance Rule" according to ILAC-G8:09/2019.

#### **Quantitative Analysis**

Quantitative values are compared with the distribution of concentration of the reference samples in the Honey-Profiling Database, for the same type of honey. Deviations to the reference range can be linked with adulterations or with specificities of the honey (e.g. untypical floral/ plant sources or production regions). For this reason, an expert interpretation is suggested in case of deviations.

All conformity statements in this section are based on the "Binary Statement for Simple Acceptance Rule" according to ILAC-G8:09/2019.

#### **Univariate and Multivariate Verification Models**

Verification models are non-targeted analyses comparing the whole NMR-Profile of a specific sample with one corresponding group of reference spectra (within the Honey-Profiling Database). All spectral data points are taken into account irrespective of whether the signals are caused by already identified molecules or not.

In the univariate analysis, the NMR spectrum is checked for any unusual low or high signal intensities, while taking into account the natural variability of a respective reference group. The chemical shifts (positions of the signals in the spectra) of the deviating signals are indicated. A guideline gives a list of possible molecules with their chemical shifts that could be responsible for the deviations.

The multivariate models take into account the relation between different signals in the NMR spectrum.

Deviations to the group of reference spectra can be linked with adulterations or specificities of the honey (e.g. untypical floral/ plant sources or production regions). For this reason, an expert interpretation is suggested in case of deviations.

All conformity statements in this section are based on the "Binary Statement for Simple Acceptance Rule" according to ILAC-G8:09/2019.

#### FAQ - Frequently asked Questions

For more information please visit our website and read our FAQ at

http://www.bruker.com/en/products-and-solutions/mr/nmr-food-solutions/honey-profiling/honey-profiling-faq.html

#### **END OF REPORT**